


SECOND EDITION

CELLULAR AND MOLECULAR IMMUNOLOGY



ABUL K. ABBAS

PROFESSOR OF MEDICINE

ANDREW H. LICHTMAN

PROFESSOR OF MEDICINE

JORDAN S. POBER

PROFESSOR OF MEDICINE




Exhibit A

SECOND EDITION

CELLULAR AND MOLECULAR IMMUNOLOGY

ABUL K. ABBAS, M.B.B.S.

Professor of Pathology
Harvard Medical School and Brigham and Women's Hospital
Boston, Massachusetts

ANDREW H. LICHTMAN, M.D., Ph.D.

Assistant Professor of Pathology
Harvard Medical School and Brigham and Women's Hospital
Boston, Massachusetts

JORDAN S. POBER, M.D., Ph.D.

Professor of Pathology, Immunobiology, and Biology
Yale University School of Medicine
New Haven, Connecticut

W.B. SAUNDERS COMPANY
A Division of Harcourt Brace & Company

Philadelphia London Toronto Montreal Sydney Tokyo

W.B. SAUNDERS COMPANY
A Division of Harcourt Brace & Company

The Curtis Center
Independence Square West
Philadelphia, Pennsylvania 19106

Library of Congress Cataloging-in-Publication Data

Abbas, Abul K.

Cellular and molecular immunology / Abul K. Abbas, Jordan S. Pober,
Andrew H. Lichtman. — 2nd ed.

p. cm.

Includes bibliographical references and index.

ISBN 0-7216-5505-X

1. Cellular immunity. 2. Molecular immunology. I. Lichtman,
Andrew H. II. Pober, Jordan S. III. Title. IV. Title: Cellular
and molecular immunology.

[DNLM: 1. Immunity, Cellular. 2. Lymphocytes—immunology. QW
568 A122c 1994]

QR185.5A23 1994

616.07'9—dc20

DNLM/DLC

93-50216

Cellular and Molecular Immunology, 2nd edition

ISBN 0-7216-5505-X

I.E. ISBN 0-7216-5290-5

Copyright © 1994, 1991 by W.B. Saunders Company

All rights reserved. No part of this publication may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopy, recording, or any information storage and retrieval system, without permission in writing from the publisher.

Printed in the United States of America.

Last digit is the print number: 9 8 7 6 5 4 3 2

ter, many of the effector functions of immunoglobulins are mediated by the Fc portions of the molecule.

Different results are obtained when the proteolytic enzyme pepsin is used instead of papain to cleave rabbit IgG molecules (Fig. 3-8). In this case, under limiting conditions of enzyme concentrations and time, proteolysis is restricted to the carboxy terminus of the hinge region near the C₂ domain such that the antigen-binding fragment of IgG retains the hinge and the interchain disulfide bonds. Fab fragments containing the heavy chain hinge are called Fab'; when the interchain disulfide bonds are intact, the two Fab' fragments remain associated in a form called F(ab')₂. The Fc fragment is often extensively degraded and does not survive proteolysis by pepsin. Fab and F(ab')₂ are often useful as experimental tools because they can bind to antigens without activating Fc-dependent effector mechanisms.

These proteolysis experiments are not readily extended to other antibody isotypes such as IgM. In fact, they are not even applicable to all IgG molecules in many species other than rabbit. However, the basic organization of the Ig molecule that Porter deduced from his studies of rabbit IgG is common to all Ig molecules of all isotypes and of all species. These features may be summarized as follows:

1. Each V_LV_H pairing forms an independent antigen-binding site. Thus, all monomeric IgG molecules have two separate antigen-binding sites, and secreted pentameric IgM molecules have ten separate antigen-binding sites (see Figs. 3-2 and 3-6).
2. The structure of the hinge region (or lack of one in certain isotypes) sterically determines how many binding sites of a single antibody molecule can simultaneously interact with antigen molecules, e.g., on a cell surface.
3. The Fc portion of an antibody molecule is spatially distinct from and functions independently of the antigen-binding site formed by the Fab regions. Since Fc regions activate immune effector functions, the kinds of effector functions activated by a particular Ig molecule are largely independent of the specificity for antigen and instead depend primarily on the isotype of the antibody.

ANTIBODY BINDING OF ANTIGENS

In the preceding sections, we have developed a general description of the structure of antibody molecules. Now we will turn to a more detailed discussion of the structural basis and physicochemical characteristics of antigen binding.

Structural Aspects of Biologic Antigens


An **antigen** can be defined as any substance that may be specifically bound by an antibody molecule. This differs from the original (historical) definition of

antigen as a molecule that generates an antibody. We now know that almost every kind of biologic molecule, including simple intermediary metabolites, sugars, lipids, autacoids, and hormones as well as macromolecules such as complex carbohydrates, phospholipids, nucleic acids, and proteins, can serve as antigens. However, only macromolecules can initiate lymphocyte activation necessary for an antibody response. Molecules that generate immune responses are called **immunogens**. (Although technically less precise, the more inclusive term "antigens" is still commonly used to refer to "immunogens.") In order to generate antibodies specific for small molecules, immunologists commonly attach such small molecules to macromolecules before immunization. In this system, the small molecule is called a **hapten** and the macromolecule, usually a foreign protein, is called a **carrier**. The hapten-carrier complex, unlike free hapten, can act as an immunogen.

In general, macromolecules are much bigger than the antigen-binding region of an antibody molecule. Therefore, an antibody binds to only a specific portion of the macromolecule, called a **determinant**, or **epitope**. These two words are synonymous and are used interchangeably throughout the book. A hapten may be thought of as an exogenous determinant that is attached to a macromolecule.

Macromolecules typically contain multiple determinants, each of which, by definition, can be bound by an antibody. In some cases, the determinants are spatially well separated, and two individual antibody molecules can be bound to the same antigen molecule without influencing each other; such determinants are said to be non-overlapping. In other cases, the first antibody bound to an antigen may sterically interfere with the binding of the second, and the determinants of the antigen are said to be overlapping. In rarer cases, binding of the first antibody may cause a conformational change in the structure of the antigen, influencing the binding of the second antibody by means other than steric hindrance. Such interactions are called allosteric effects.

In the case of phospholipids or of complex carbohydrates, the antigenic determinants are entirely a function of the covalent structure of the macromolecule. However, in the case of nucleic acids, and even more so in the case of proteins, the non-covalent folding of the macromolecule may also contribute to the formation of determinants. In proteins, epitopes formed by adjacent amino acid residues in the covalent sequence are called **linear determinants** (Fig. 3-9). It is estimated that, in a protein antigen, the size of the linear determinant that forms contacts with specific antibody is about six amino acids long. Linear determinants may be accessible to antibodies in the native folded protein if they appear on the surface or in a region of extended conformation. More often, linear determinants may be inaccessible in the native conformation and appear only when the protein is denatured. In contrast, **conformational determinants** are formed by amino acid residues from separated portions of the linear amino acid sequence that are spatially juxtaposed only upon folding (Fig. 3-9). In theory, denatured proteins could transiently give rise to conformational



1999 Catalog




Exhibit B

R&D

Antibodies

R&D Systems' polyclonal and monoclonal antibodies to cytokines, receptors and related molecules are raised against purified proteins and affinity-purified using either protein A, G, or antigen columns. Each antibody has been tested for neutralization of bioactivity as well as for use in direct ELISA and western blot applications. Many antibodies have been labeled and tested for use in flow cytometry applications and for building ELISAs. For more information on matched antibody pairs for ELISA, refer to the section on ELISA development.

Antigen affinity purified polyclonal antibodies are at least 10-fold more potent than protein G affinity purified antibodies. The use of analyte-specific affinity chromatography almost eliminates non-specific immunoglobulin. The polyclonal antibody pool recognizes more than one antigenic epitope giving a stronger staining signal than monoclonal antibodies. Many affinity purified antibodies have been tested by scientists world-wide for cytokine-specific immunostaining.

Antibodies To Cytokines & Related Molecules

6CKine 134	CD40 Ligand 154	Fractalkine 177	IL-6 R 213-214	MIP-1 α 241-242	SCF 261-262
6X-Histidine 134	CD44 154	G-CSF 177-178	IL-7 214-215	MIP-1 β 242	SCF R 262
Activin RI 134	CD44H 155	G-CSF R 178	IL-7 R 215	MIP-1 β 243-244	SDF-1 263
Activin RIIA 135	CD44v3 155	GCP-2 179	IL-8 216	MIP-II 244	SDF-1 β 263
Activin RIIIB 135	CD44v4/5 155	GDNF 179-180	IL-9 217	MIP-1 γ 244	E-/P-Selectin 263
ANG 135-136	CD44v6 155	GFR α 1 180	IL-9 R 218	MIP-1 δ 245	E-Selectin 263-264
Annexin V 136	CINC-1 156	GM-CSF 180-182	IL-10 218-219	MIP-2 245	L-Selectin 264
AR 136-137	CINC-2 α 156	GPIIb/GPIIIa 182	IL-10 R 220	MIP-3 α 246	P-Selectin 265
B7-1 137	CINC-2 β 156	gp130 182-183	IL-11 220-221	MIP-3 β 246	Shh 265
B7-2 137-138	CNTF 156-158	GRO 183	IL-11 R α 221	MK 246	SLPI 266
Bad 138	CNTF R α 158	GRO α 183-184	IL-12 222-223	MMP-1 247	Stat 1p91 266
Bag-1 138	CPP32 159	HB-EGF 184	IL-12 p40 223	MMP-2 247	Stat 2 266
BAK 139	CRG-2 159	HCC-1 184	IL-12 p70 223	MMP-3 248	Stat3 266
Bax 139-140	CTLA-4 159	HCC-4 185	IL-13 224-225	MMP-7 248	Stat 4 267
Bcl-2 140	CXCR-1 160	HGF 185	IL-15 225	MMP-9 248	Stat 5A 267
Bcl-x 141	CXCR-2 160	HRG- α 186	IL-16 226	MMP-12 249	Stat 5B 267
Bcl-x ℓ 141	CXCR-3 161	HRG- β 1 186	IL-17 226-227	MPIF-1 249	Stat 6 268
BDNF 141-142	CXCR-4 161-163	I-309 186	IL-18 227-228	MSP 249	TARC 268
BL-CAM 142	CXCR-5 163-164	clAP-1 (HIAP-2) 187	IP-10 228	MSP R 250	TECK 268-269
BLC/BCA-1 142-143	Cytochrome c 164	clAP-2 (HIAP-1) 187	Jak-1 229	NAP-2 250	TGF- α 269
BMP-2 143	EGF 165	ICAM-1 187-188	KC 229	NCAM 250	TGF- β 269
BMP-5 143	ENA-78 166	ICAM-3 188-189	Leptin 229-230	β -NGF 250-251	TGF- β 1 270
BMPR-IA 143	Eotaxin 167	IFN- γ 189-191	Leptin R 230-231	NGF R 251	LAP 270-271
BTC 144	Eotaxin-2 (MPIF-2) 168	IGF-I 191-192	LFA-1 231	eNOS 252	TGF- β 2 271-272
C10 144-145	ephrin-A2 168	IGF-I R 192	LFA-1 α 231	NT-3 252	TGF- β 3 272-273
E-Cadherin 145-146	ephrin-A5 168	IGF-II 192	LFA-1 β 232	NT-4 252-253	TGF- β 5 273
N-Cadherin 146	Epo 169	IL-1 α 193-195	LIF 232-233	OSM 253-254	LTBP-1 273
P-Cadherin 146	Epo R 169	IL-1 β 195-197	LIF R 233	Osteopontin 254	TGF- β RII 274
P-, E-, N-Cadherin 147	Fas 170	IL-1 RI 197-198	Lymphotactin 234	p150,95 α chain 254	TGF- β RIII 274
CCR-1 147	FGF acidic 170-171	IL-1 RII 198-199	M-CSF 234-235	PARC 254	Tie 1 274
CCR-2 147-148	FGF basic 171	IL-1 α R 199-200	M-CSF R 235	PARP 255	TNF- α 274-277
CCR-3 148	FGF-4 172	IL-2 200-202	Mac-1 α 235	PD-ECGF 255	TNF- β 277-278
CCR-5 148-150	FGF-5 173	IL-2 R α 202-203	MARC 236	PDGF 256	TNF RI 278-279
CCR-6 151	FGF-6 173	IL-2 R β 203	MCP-1 236-237	PDGF-AA 256	TNF RII 279-280
CD3 ϵ 151	FGF-7 174	IL-2 R γ 204	MCP-1/JE 237	PDGF-BB 257	Tpo 280-281
CD4 152	FGF-8 174	IL-3 204-206	MCP-2 237-238	PDGF R α 257	TrkC 282
CD14 152	FGF-8b 174	IL-3 R α 206	MCP-3 238-239	PDGF R β 258	uPAR 282
CD15 153	FGF-9 175	IL-4 206-208	MCP-4 239	PIGF 258	VCAM-1 282-283
CD27 153	FGF-10 175	IL-4 R 209	MCP-5 239	PIGF-2 259	VEGF 283-284
CD28 153	Flt-1 175	IL-5 209-211	MDC 239	PTN 259	VLA-4 284
CD31 154	Flt-3 Ligand 176	IL-5 R α 211	MIF 240	RANTES 259-260	WNT-4 284
CD40 154	Flt-4 177	IL-6 211-213	MIG 240-241	Ret 261	

anti-Bcl-x
polyclonal antibody

Catalog #	AF800
Size/Price	50 µg/\$190
Form	Antigen affinity-purified, 0.2 µm filtered solution in PBS and NaN ₃ , lyophilized
Type	Mouse/human Bcl-x specific rabbit IgG
Immunogen	Synthetic peptide corresponding to amino acids 49 - 68 of the Bcl-x sequence
Specificity	Selected for its ability to react with human and mouse Bcl-x.
Western Blot	An antibody concentration of 0.3 µg/mL is recommended.
Immuno-precipitation	3 µg of antibody per immunoprecipitation of Bcl-x from 7 x 10 ⁵ A431 cells is recommended.

anti-Bcl-x_L
monoclonal antibody

Catalog #	2300-MC-100
Size/Price	100 µg/\$230
Form	0.2 µm filtered solution in PBS, liquid
Type	Mouse IgG _{2a} , clone # YTH-2H12
Immunogen	Synthetic peptide corresponding to amino acids 3 to 14 of the human Bcl-x _L sequence. This epitope is also present in Bcl-X _s , a variant of an alternative RNA splicing.
Specificity	Selected for its ability to recognize human, mouse and rat Bcl-x _L .
Applications	See product insert for performance specifications.

anti-human BDNF
polyclonal antibody

Catalog #	AF248
Size/Price	100 µg/\$340
Form	Antigen affinity-purified, 0.2 µm filtered solution in PBS, lyophilized
Type	Human BDNF specific chicken IgG
Immunogen	Sf21-expressed recombinant human BDNF
Specificity	Selected for its ability to recognize rhBDNF.
Applications	See product insert for performance specifications.

anti-human BDNF
monoclonal antibody

Catalog #	MAB248
Size/Price	500 µg/\$280
Form	0.2 µm filtered solution in PBS, lyophilized
Type	Mouse IgG ₁ , clone # 35928.11
Immunogen	Sf21-expressed recombinant human BDNF
Specificity	Selected for its ability to neutralize the bioactivity of rhBDNF.
Neutralization	5 - 15 µg/mL will neutralize 50% of the bioactivity due to 2.5 ng/mL of rhBDNF.
ELISA	0.5 - 1 µg/mL will detect <12.5 ng/well of rhBDNF.
Western Blot	1 - 2 µg/mL, with the appropriate secondary reagents, will allow visualization of approximately 300 ng/lane of rhBDNF under both non-reducing and reducing conditions.

Antibodies
to
Cytokines
& Related
MoleculesBcl-x
Bcl-x_L
BDNF**For Technical Service Or To Place An Order**

USA & Canada: 800-343-7475 (Tele) UK & Europe: +44 (0)1235 551100 (Tele)

612-627-0424 (Fax)

+44 (0)1235 533420 (Fax)

www.rndsystems.com

Copied from 10729653 on 09/08/2004

anti-CPP32 polyclonal antibody

Catalog #	AF-605-NA
Size/Price	100 µg/\$230
Form	Antigen affinity-purified, 0.2 µm filtered solution in PBS and NaN ₃ , lyophilized
Type	CPP32 specific goat IgG
Immunogen	Recombinant human CPP32 zymogen
Specificity	Selected for its ability to react with intact human CPP32 and with CPP32 fragments of ~20, 18, and 16 kDa that are produced during apoptosis.
Western Blot	0.5 µg/mL, with the appropriate secondary reagents, is recommended.
Immuno-precipitation	4 µg of antibody per immunoprecipitation of human CPP32 from 1 - 2 x 10 ⁶ cells is recommended.

anti-mouse CRG-2/IP-10 polyclonal antibody

Catalog #	AF-466-NA
Size/Price	100 µg/\$340
Form	Antigen affinity-purified, 0.2 µm filtered solution in PBS, lyophilized
Type	Mouse CRG-2 specific goat IgG
Immunogen	<i>E. coli</i> -expressed recombinant mouse CRG-2
Specificity	Selected for its ability to neutralize the bioactivity of rmCRG-2. In direct ELISA and western blot applications, this antibody shows <10% cross-reactivity with rhIP-10.
Neutralization	4 - 12 µg/mL will neutralize 50% of the bioactivity due to 200 ng/mL of rmCRG-2.
ELISA	0.5 - 1 µg/mL will detect <0.16 ng/well of rmCRG-2.
Western Blot	0.1 - 0.2 µg/mL, with the appropriate secondary reagents, will allow visualization of approximately 5 ng/lane of rmCRG-2 under both non-reducing and reducing conditions.
Also Available Biotinylated (Catalog # BAF466, Size 50 µg, Price \$365)	

anti-human CTLA-4 polyclonal antibody

Catalog #	AF-386-PB
Size/Price	100 µg/\$340
Form	Antigen affinity-purified, 0.2 µm filtered solution in PBS, lyophilized
Type	Human CTLA-4 specific goat IgG
Immunogen	Sf21-expressed recombinant human CTLA-4, extracellular domain
Specificity	Selected for its ability to detect rhCTLA-4 in direct ELISA and western blot applications.
ELISA	0.5 - 1 µg/mL will detect <0.16 ng/well of rhCTLA-4.
Western Blot	0.1 - 0.2 µg/mL, with the appropriate secondary reagents, will allow visualization of approximately 5 ng/lane of rhCTLA-4 under both non-reducing and reducing conditions.

Also Available Biotinylated
(Catalog # BAF386, Size 50 µg, Price \$365)

anti-mouse CTLA-4 polyclonal antibody

Catalog #	AF476
Size/Price	100 µg/\$340
Form	Antigen affinity-purified, 0.2 µm filtered solution in PBS, lyophilized
Type	Mouse CTLA-4 specific goat IgG
Immunogen	NSO-expressed recombinant mouse CTLA-4, extracellular domain
Specificity	Selected for its ability to recognize rmCTLA-4.
Applications	See product insert for performance specifications.

Also Available Biotinylated
(Catalog # BAF476, Size 50 µg, Price \$365)



Antibodies
to
Cytokines
& Related
Molecules

CPP32
CRG-2
CTLA-4

For Technical Service Or To Place An Order

USA & Canada: 800-343-7475 (Tele) UK & Europe: +44 (0)1235 551100 (Tele)

612-627-0424 (Fax) +44 (0)1235 533420 (Fax)



www.rndsystems.com

Copied from 10729653 on 09/08/2004

**labeled anti-human CXCR-5 (BLR-1)
monoclonal antibody**

Catalog #	FAB190B	FAB190P
Size/Price	100 Tests/\$205	100 Tests/\$230
Label	Biotin	Phycoerythrin
Form	1 mL in a HEPES buffered saline containing 0.5% BSA and 0.1% NaN ₃ .	
Type	Mouse IgG _{2b} , clone # 51505.111	
Immunogen	CXCR-5 transfected NSO mouse myeloma cells	
Specificity	Selected for its ability to react by FACS analysis with human CXCR-5 transfectants. This antibody does not cross-react with CXCR-2, CXCR-3, and CXCR-4 transfectants.	
Flow Cytometry	10 µL, with the appropriate secondary reagents, can be used to stain 1x10 ⁵ cells for flow cytometry applications.	

**anti-Cytochrome c
monoclonal antibody**

Catalog #	MAB897
Size/Price	100 µg/\$200
Form	0.2 µm filtered solution in PBS, lyophilized
Type	Mouse IgG _{2b} , clone # 7H8.2C12
Immunogen	Pigeon Cytochrome c
Specificity	Selected for its ability to recognize human or mouse Cytochrome c.
Applications	See product insert for performance specifications.

**anti-human Cytochrome c
monoclonal antibody**

Catalog #	MAB898
Size/Price	100 µg/\$200
Form	0.2 µm filtered solution in PBS, lyophilized
Type	Mouse IgG _{2a} , clone # 2B5F8
Immunogen	Equine Cytochrome c
Specificity	Selected for its ability to recognize human Cytochrome c.
Applications	See product insert for performance specifications.

**anti-Cytochrome c
monoclonal antibody**

Catalog #	MAB899
Size/Price	100 µg/\$200
Form	0.2 µm filtered solution in PBS, lyophilized
Type	Mouse IgG _{2a} , clone # 2.7D5
Immunogen	Pigeon Cytochrome c
Specificity	Selected for its ability to recognize human or mouse Cytochrome c.
Applications	See product insert for performance specifications.

Antibodies
to
Cytokines
& Related
Molecules

CXCR-5
Cytochrome c

These products are for research use only, not for use in diagnostic procedures.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☒ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.